

Assignment III, PHYS 112 (General Physics II)
Fall 2020

Due via pdf upload to OAKS prior to Friday, September 11th at 10:00 AM

General instructions:


For this, and all other homework assignments, please turn in your solutions with all supporting work; answers without supporting work will not earn credit. You do not need to upload the sheet with the questions on it, but please clearly number your problems and circle or box your final answers. I encourage you to collaborate with classmates to discuss how to approach a particular question, but the mathematical steps to generate your final answer on your submitted work should be your own. If I see the same simple mistake on multiple homework assignments, I will take off more points for that error than I normally would. Please include *words* in your answers. When you get answer keys back from me, you'll see that there are explanations, ideas, commentary, and thought processes included – not just a set of equations one after another. Finally, please ensure that all numerical answers have units.

1. Let the following charges exist:

$$\begin{aligned} q_1 &= 970 \text{ nC} & \text{at} & \quad (2.0 \text{ m})\hat{i} - (1.5 \text{ m})\hat{j} \\ q_2 &= -1.30 \text{ } \mu\text{C} & \text{at} & \quad (2.0 \text{ m})\hat{i} + (0.0 \text{ m})\hat{j} \\ q_3 &= 1.80 \text{ } \mu\text{C} & \text{at} & \quad (-0.3 \text{ m})\hat{i} - (1.0 \text{ m})\hat{j} \end{aligned}$$

- a) If q_4 had charge $-2.1 \text{ } \mu\text{C}$ and was at $1.5 \text{ m}\hat{i} - 1.0 \text{ m}\hat{j}$, what would the force on this charge be due to the presence of q_1 , q_2 , and q_3 ? (You may assume q_1 , q_2 , and q_3 remain stationary). Remember, force is a vector quantity.
 - b) If q_4 had charge $+380 \text{ nC}$ and it was at $1.5 \text{ m}\hat{i} - 1.0 \text{ m}\hat{j}$ (the same position as part (a)), what would the force on this charge q_4 be due to the presence of q_1 , q_2 , and q_3 ? Again – you may assume that the other charges are stationary. (There may be a shortcut to this one if you've already done part (a)).
2. A ping-pong ball has a mass of about 2.7 grams. Let us say I was able (somehow) to put a total net charge of -34.6 nC on its surface. What minimum strength (and direction) of electric field would be required to levitate the ping-pong ball (near the surface of the Earth). (When you report the direction, report it as “up” or “down” or “East” or “West” or “North” or “South”; don't use \hat{i} , \hat{j} , \hat{k} unless it is obvious to me how that relates back to the Earth).

3. A Helium nucleus is made up of two protons and two neutrons. (For this problem, assume a neutron has the same mass as a proton, but does not have any charge). Assume that this Helium nucleus is placed in a uniform electric field of $(1.35 \times 10^5 \text{ N/C})\hat{i}$.
- Assume the Helium nucleus is released from rest at the origin at time $t = 0$. Find the speed of the Helium nucleus after it travels 1.00 cm. [Hint – constant electric field means constant force. Hopefully you remember some stuff from 111].
 - The speed of light in a vacuum is $3 \times 10^8 \text{ m/s}$. No matter can actually move that fast (we'll find out why later this semester). However, if matter *could* travel that fast (and all the results you learned in PHYS 111 hold), where would the nucleus be when it reaches the speed of light?
 - How long does it take the Helium nucleus to reach the location identified in part (b) above?
4. The Earth – near its surface – has its own electric field of approximately 150 N/C (pointing towards the center of the Earth).
- What is the approximate net charge of the Earth? (Treat the Earth as a point-mass and point-charge at its center).
 - What magnitude and sign of net charge would a person have to obtain in order to levitate due to the superposition of gravitational and electric forces?
 - Let's say two people each have accumulated the net charge calculated in part (b). What would their force of repulsion be if they were 1 meter apart?
5. Examine the figure below, corresponding to a fixed charge configuration. (In other words, q_1 , q_2 , q_3 , and q_4 are not able to move). In this problem, leave your answers in terms of Q , d , and fundamental constants.
- If $q_1 = q_2 = q_3 = q_4 = Q$, what is the magnitude of the electric field at point P?
 - If $q_1 = q_2 = Q$ and $q_3 = q_4 = -Q$, what is the magnitude of the electric field at point P?
 - If $q_1 = q_4 = Q$ and $q_2 = q_3 = -Q$, what is the magnitude of the electric field at point P?
 - If $q_1 = q_2 = q_3 = Q$ and $q_4 = -Q$, what is the magnitude of the electric field at point P?



6. If you didn't know, inkjet printers work through an electromechanical interaction. What happens is that drops of ink are "spit" at the paper from a nozzle. The ink drops each have a mass of about 150 ng and travel toward the paper at about 20 m/s. During their travel, the drops go through a "charging electrode" that assigns each drop some positive charge q by removing some electrons. The drops then pass between parallel plates that are about 2 cm long where a uniform electric field of magnitude about 80 kN/C is applied. (The plates are oriented similar to what is shown in the figure below so that they can generate some deflection to the droplet's straight-line path). If a drop needs to be deflected 0.4 mm by the time it reaches the end of the deflection plates, how many electrons were removed from the drop?

