

**Assignment IX, PHYS 111 (General Physics I)**  
**Fall 2021**  
**Due 11/4/21**

Just as a reminder – in each homework assignment, I will list suggested homework problems out of the book. These are worth practicing – some may even appear on exams verbatim – but since they are in the text, finding answers on-line should be straightforward and these textbook problems will not be graded. I suggest you do them – many of them will be easier than the graded homework and they would be a good thing to tackle in your SI sessions to get comfortable with the content.

After the suggested book problems, I will give a list of problems that I myself wrote. *SOME* of these problems will be graded, but you won't know which ahead of time. The ones that I grade will be the same for everyone in the class.

I will supply you with an answer key to all of the problems that I wrote – even the ones that I did not grade.

As always, please legibly write (or type) your answers on separate paper. Incorrect answers with no work will earn nearly no credit, and consistent correct answers with no work are suspicious – many of these problems your professor can't do in his head, so it is unusual if you can. Please show all relevant work.

To help with this homework, you should read the associated sections of your text and watch the videos associated with the lectures on the course webpage: [http://larsenml.people.cofc.edu/phys111\\_fall21.html](http://larsenml.people.cofc.edu/phys111_fall21.html).

## **(Ungraded) suggested textbook practice problems**

(All problems are odd problems (that have answers in the back of the book) out of Halliday, Resnick, and Walker, 10th Ed.) Some problems from previous chapters that are related to recent lecture content have been included.

Chapter 10:


Problems: 45, 47, 53, 71, 75, 89

Chapter 11:

Problems: 3, 5, 7, 11, 19, 23, 27, 33, 37, 39, 45, 49, 51, 79, 83, 85


## Graded Homework Problems

- Let mass  $m_1=2\text{kg}$  be located at  $2m \hat{j}$  and  $m_2 = 1\text{kg}$  be located at  $3m \hat{i}$ . The masses are connected to each other and to the axis of rotation by massless, rigid rods. Find the moments of inertia for this mass configuration with respect to the following rotational axes. (Your answer should be different for each axis).
  - The  $x$ -axis
  - The  $y$ -axis
  - The  $z$ -axis
  - The straight-line axis that goes through both  $m_1$  and  $m_2$ .
- A standard, hinged door can be treated as a 2-dimensional plate with dimensions  $h$  and  $w$  (for height and width, respectively). The moment of inertia of a flat rectangular plate like this with respect to its hinged side would be  $I = \frac{1}{3}mw^2$ . Let us say that, for a real door, this moment of inertia ends up being equal to  $25 \text{ kg m}^2$ , and the door has a width of 1.3 meters.
  - What is the mass of the door? (They're not all hard!)
  - Neglecting friction, what steady force – applied at its outer edge and perpendicular to the surface of the door – can move the door from rest through an angle of  $\pi/2$  radians in 1.4 seconds?
  - If the force was applied to a position 0.5 meters from the hinges (but still perpendicular to the surface of the door), what steady force would have needed to be applied in order to move the door from rest through an angle of  $\pi/2$  radians in 1.4 seconds?
- The moment of inertia of a compound object is equal to the sum of the moments of inertia of each piece. The moment of inertia of a uniform disk about its center is  $\frac{1}{2}MR^2$  where  $M$  is the mass of the disk and  $R$  is the radius of the disk. Two uniform metal disks with masses  $m_1$  and  $m_2$  and radii  $R_1$  and  $R_2$  (respectively) are welded together and mounted on a frictionless axis through their common center as shown in the figure below.
  - What is the total moment of inertia of the two disks with respect to the axis shown? (Leave your answer in terms of  $m_1$ ,  $m_2$ ,  $R_1$ , and  $R_2$ ).
  - Let  $R_1 = 7.00 \text{ cm}$  and  $R_2 = 11.0 \text{ cm}$ . Let  $m_1 = 1.35 \text{ kg}$  and  $m_2 = 4.23 \text{ kg}$ . A massless, inextensible string is wrapped around the edge of the smaller disk and a  $1.50 \text{ kg}$  block is suspended from the free end of the string. If the block is released from rest a distance  $10.00 \text{ m}$  above the floor, what is the speed of the block just before it strikes the floor?
  - Repeat your calculation in part (b) above if the string is wrapped around the edge of the larger disk instead.




4. Consider the system displayed below. The mass distribution in both wheels is not uniform, but rather it is weighted towards the outside. This means that the moment of inertia of each of these wheels is not  $\frac{1}{2}MR^2$  like it would be for a uniform disk, but rather it is  $\frac{3}{4}MR^2$ . The wheels are welded together and – as shown – a 50 N force is applied from a rope to the right connected to the smaller wheel. The mass and radius of the smaller wheel are 2.5 kg and 35 cm, respectively. The mass and radius of the larger wheel are 5.2 kg and 45 cm, respectively.

- How large must the force on the rope connected to the larger wheel be to prevent the system from rotating?
- How large must the force on the rope connected to the larger wheel be if the system goes from rest to completing 5 full rotations counter-clockwise in 18.0 seconds?
- Continuing on from part (b) above, what would be the angular momentum of the whole system after 18.0 seconds if the force calculated in part (b) is applied to the rope connected to the larger wheel?



5. Examine the figure below. The pulley has moment of inertia  $I$ .

- Find  $m_2$  in terms of  $a$ ,  $m_1$ ,  $r_1$ ,  $r_2$ ,  $I$ , and  $g$  if  $m_2$  accelerates downwards with acceleration  $a$ .
- Find  $m_2$  in terms of  $a$ ,  $m_1$ ,  $r_1$ ,  $r_2$ ,  $I$ , and  $g$  if  $m_2$  accelerates upwards with acceleration  $a$ .



6. A force of  $(30\hat{i} - 25\hat{j} + 10\hat{k})\text{N}$  is applied at  $\vec{r} = (4\hat{i} + 3\hat{j} - 2\hat{k})\text{m}$ . What is the magnitude and direction of the torque of this force about the origin?

7. A force of 35N at  $\pi/7$  radians with respect to the  $+x$  axis is applied to the edge of a large wheel with radius 7 m centered at the origin. The force is applied at the position on the edge of the wheel an angle of  $6\pi/5$  radians with respect to the  $+x$  axis. What is the magnitude of the torque from this force?
8. Under certain circumstances, a star can collapse into an extremely dense object called a neutron star. (1 tablespoon full of a neutron star has approximately the same mass as the whole Earth). Let's talk about a star that has an initial radius  $9 \times 10^8$  m and collapsed down into a neutron star that has a radius of about  $2 \times 10^4$  m. You may assume that the mass of each star is uniformly distributed throughout its volume.
- a) If no mass is lost in this process, what is the rotational period of the neutron star if the original star had a rotational period of 2 weeks?
- b) Find the ratio  $\frac{v_{\text{rim}}^{\text{neutron star}}}{v_{\text{rim}}^{\text{original star}}}$ , where  $v_{\text{rim}}$  corresponds to the velocity of a piece of the star at the surface and on the equator of the rotating star.